



DR MAHA OTHMAN (Orcid ID : 0000-0001-7562-203X)

Article type : Letter to the Editor

Exploring possible mechanisms for COVID-19 induced thrombocytopenia: Unanswered Questions

Ariunzaya Amgalan,¹ Maha Othman^{2,3}

1. Georgetown University School of Medicine, Washington, DC, USA.
2. Department of Biomedical and Molecular Sciences, School of Medicine, Queen's University, Kingston, Canada.
3. School of Baccalaureate Nursing, St. Lawrence College, Kingston, Canada.

Corresponding author:

Maha Othman MD PhD

Department of Biomedical and Molecular Sciences

School of Medicine, Queen's University

Kingston, Ontario, Canada

e-mail: othman@queensu.ca

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/JTH.14832](https://doi.org/10.1111/JTH.14832)

This article is protected by copyright. All rights reserved

Dear Drs. Lillicrap and Morrissey,

We read with enthusiasm the recent article by Tang et al. entitled “Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia,” which is a retrospective analysis of the coagulation profile of 183 patients admitted with pneumonia secondary

to the 2019-nCoV (COVID-19).¹ This article deals with a timely topic and brings up important points that need to be interrogated in future studies.

Tang et al. reported 11.5% mortality in patients with COVID-19 pneumonia and noted that 71.4% of these deaths had abnormal coagulation profiles consistent with disseminated intravascular coagulation (DIC) based on criteria from the International Society on Thrombosis and Haemostasis (ISTH).¹ Compared to survivors of COVID-19 pneumonia, the non-survivors had significantly higher D-dimer and fibrin degradation product levels, and longer prothrombin time and activated partial thromboplastin time.¹ Consistent with the findings of Tang et al., a meta-analysis of 1779 patients with COVID-19 found that a low platelet count was associated with over fivefold increased risk of severe disease (OR, 5.1; 95% CI, 1.8-14.6) and an even lower platelet count was associated with mortality in those patients.²

It is interesting that 57.1% of the non-survivors in the study by Tang et al., (12 out of 21) exhibited thrombocytopenia (7 (33.3%) had $50\sim 100 \times 10^9/L$ and 5 (23.8%) had $<50 \times 10^9/L$ platelets).¹ While thrombocytopenia is a key diagnostic component in DIC, the data in this study raises a question of whether thrombocytopenia in COVID-19 is a part of sepsis-induced DIC and/or a direct platelet-viral interaction and if so, is this interaction beneficial for the host or for the virus and what are the possible mechanisms. These questions are worth exploring further in research studies.

The pathophysiology of DIC is complex, involving combined activation of the vascular endothelium, platelets, and leukocytes, and results in thrombin generation and widespread fibrin deposition.³ There is multiple evidence that platelets interact directly with viral pathogens via pathogen recognition receptors (PRR) and this is part of their role in fighting infections.⁴ Platelets express PRRs such as C-type lectin receptors and toll-like receptors TLR-3, -7, and -9, which are reported to have a role in platelet-viral infections.⁴ This interaction subsequently leads to platelet activation and release of molecules with antiviral activity including kinocidins and other microbicidal peptides.⁴ Activated platelets are known to produce reactive oxygen species and can also interact with and activate other immune cells such as leukocytes which can also fight and clear viral pathogens.⁴ Activated platelets

are cleared from the circulation by the reticuloendothelial system.^{4,5} Thrombocytopenia has been reported with several viral infections such as HIV, Influenza, Hepatitis B and C viruses, CMV, EBV and Rhinovirus.⁵

Viruses can directly interact with platelets, thereby altering their count and/or function.^{5,6}

Platelet specific receptors known to interact with viruses are reported.⁶ Various mechanisms were explained and are dependent on the virus type. These include: triggering systemic inflammation and clearance of activated platelets via splenic/liver macrophages and/or phagocytosis by neutrophils such as the case in influenza and rhinoviruses,⁵ suppressing platelet production or enhancing platelet destruction such as the case in herpes and simian viruses,⁵ and anti-viral antibodies cross-reacting with platelet surface integrins such as the case in adenovirus (platelet integrin GPIIb/IIIa).^{5,7} Platelets were found to express the Coxsackie adenovirus receptor (CAR) and in-vitro and in-vivo studies showed direct adenovirus-platelet interactions, followed by platelet activation as evidenced by increased expression of P-selectin with subsequent formation of platelet-leukocyte aggregates and expression of leukocytes and endothelial cells' activation markers.^{8,9} Furthermore, the adenovirus-induced thrombocytopenia was found to be dependent on von Willebrand factor (vWF) since vWF-KO mice did not show thrombocytopenia when injected by the virus.⁸

For thrombocytopenia associated with severe acute respiratory syndrome-coronavirus (SARS-CoV), the most closely related virus to COVID-19, multiple possible mechanisms have been suggested including the development of autoantibodies or immune complexes mediating clearance, direct infection of hematopoietic stem/progenitor cells and the megakaryocytic lineage via CD13 or CD66a resulting in decreased production of platelets, increased thrombomodulin levels leading to pathologic activation of the coagulation pathway and consumption of platelets, and increased plasma concentrations of tissue-plasminogen activator reflecting increased fibrinolysis.^{10,11} Chen et al. found that patients with SARS had significantly lower lymphocyte and platelet counts, and significantly higher soluble vascular cell adhesion molecule-1 (sVCAM-1) and soluble Fas ligand (sFasL) levels compared to healthy patients.¹² However, it remains unclear whether the apoptosis pathway is responsible for the reduced blood cell counts or whether sFasL and sVCAM-1 reflect vascular or

tissue damage.¹² Researchers have also provided evidence that dysregulation of the urokinase pathway contributes to SARS-CoV pathogenesis.¹³ Gralinski et al. demonstrated a protective role for *Serpine1*, an inhibitor of the urokinase pathway, by showing that *Serpine1* knockout mice infected with SARS-CoV had increased fibrinolytic activity in their lungs and a decreased inflammatory response compared to wild-type mice.¹³ Citing evidence from previous literature on SARS-CoV infection, Tang et al. discusses a possibility of dysregulation of the urokinase pathway and provides evidence of fibrinolysis with COVID-19 deaths.¹

To date, it is unclear what the exact mechanism of COVID-19 induced thrombocytopenia is. We don't know if this is a direct platelet-COVID-19 virus interaction or part of the sepsis-induced DIC. Is this interaction beneficial for the host; a trial to clear the viral pathogen or for the virus; overcoming innate immunity. Is there a specific platelet ligand that COVID-19 binds to? What we do know is that COVID-19 belongs to the same lineage of coronaviruses as SARS-CoV, so it may be helpful to look at the mechanism behind SARS-CoV to guide COVID-19 studies while also keeping in mind that there are significant differences between SARS-CoV and COVID-19.¹⁴ Compared to SARS-CoV, COVID-19 spike proteins have at least 10 times greater affinity for the human angiotensin-converting enzyme 2 (ACE2), a common host cell receptor shared between the two viruses.¹⁴ Furthermore, Wrapp et al. demonstrated that three different antibodies known to bind to the SARS-CoV spike protein failed to bind to COVID-19, suggesting the uniqueness of the novel coronavirus.¹⁴

A simple algorithm for the management of COVID-19 coagulopathy has just been reported. This is only based on supportive therapy, given the currently limited knowledge of the pathophysiology of DIC in COVID-19.¹⁵ An understanding of the innate immune response and interrelated hemostatic mechanisms as critical initial response to COVID 19 infection is needed.¹⁶ Additional research is warranted to elucidate the causal and pathophysiologic mechanisms of DIC and thrombocytopenia associated with COVID-19 with insight into the innate immune response to this virus. As the COVID-19 pandemic continues to evolve, research into these matters would be of utmost importance.

Word count: 1085

Conflict of Interest Disclosure: all authors have nothing to disclose.

Author Contribution: AA gathered literature and wrote the manuscript. MO developed the plan and rationale for the letter, gathered literature and wrote the manuscript.

References:

1. Tang N, Li D, Wang X, Sun Z. Abnormal Coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost.* 2020. doi:10.1111/jth.14768
2. Lippi G, Plebani M, Michael Henry B. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A meta-analysis. *Clin Chim Acta.* March 2020. doi:10.1016/j.cca.2020.03.022
3. Gando S, Levi M, Toh CH. Disseminated intravascular coagulation. *Nat Rev Dis Prim.* 2016;2. doi:10.1038/nrdp.2016.37
4. Hottz ED, Bozza FA, Bozza PT. Platelets in immune response to virus and immunopathology of viral infections. *Front Med.* 2018;5(APR). doi:10.3389/fmed.2018.00121
5. Assinger A. Platelets and infection - An emerging role of platelets in viral infection. *Front Immunol.* 2014;5(DEC). doi:10.3389/fimmu.2014.00649
6. Seyoum M, Enawgaw B, Melku M. Human blood platelets and viruses: Defense mechanism and role in the removal of viral pathogens. *Thromb J.* 2018;16(1):16. doi:10.1186/s12959-018-0170-8
7. Gupalo E, Kuk C, Qadura M, Buriachkovskaia L, Othman M. Platelet-adenovirus vs. inert

- particles interaction: Effect on aggregation and the role of platelet membrane receptors. *Platelets*. 2013;24(5):383-391. doi:10.3109/09537104.2012.703792
8. Othman M, Labelle A, Mazzetti I, Elbatarny HS, Lillicrap D. Adenovirus-induced thrombocytopenia: The role of von Willebrand factor and P-selectin in mediating accelerated platelet clearance. *Blood*. 2007;109(7):2832-2839. doi:10.1182/blood-2006-06-032524
 9. Gupalo E, Buriachkovskaia L, Othman M. Human platelets express CAR with localization at the sites of intercellular interaction. *Viol J*. 2011;8:456. doi:10.1186/1743-422X-8-456
 10. Goeijenbier M, van Wissen M, van de Weg C, et al. Review: Viral infections and mechanisms of thrombosis and bleeding. *J Med Virol*. 2012;84(10):1680-1696. doi:10.1002/jmv.23354
 11. Yang M, Li CK, Li K, et al. Hematological findings in SARS patients and possible mechanisms (review). *Int J Mol Med*. 2004;14(2):311-315. doi:10.3892/ijmm.14.2.311
 12. Chen RF, Chang JC, Yeh WT, et al. Role of vascular cell adhesion molecules and leukocyte apoptosis in the lymphopenia and thrombocytopenia of patients with severe acute respiratory syndrome (SARS). *Microbes Infect*. 2006;8(1):122-127. doi:10.1016/j.micinf.2005.06.007
 13. Gralinski LE, Bankhead A, Jeng S, et al. Mechanisms of severe acute respiratory syndrome coronavirus-induced acute lung injury. *MBio*. 2013;4(4). doi:10.1128/mBio.00271-13
 14. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*. February 2020. doi:10.1126/science.abb2507
 15. Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. March 2020. doi:10.1111/jth.14810
 16. Lillicrap D. Disseminated intravascular coagulation in patients with 2019-nCoV pneumonia. *J Thromb Haemost*. March 2020. doi:10.1111/jth.14781